

Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: A systematic review and network meta-analysis

Lukas Schwingshackl <https://orcid.org/0000-0003-3407-7594>, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann & Heiner Boeing

To cite this article: Lukas Schwingshackl <https://orcid.org/0000-0003-3407-7594>, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann & Heiner Boeing (2019) Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: A systematic review and network meta-analysis, Critical Reviews in Food Science and Nutrition, 59:16, 2674-2687, DOI: [10.1080/10408398.2018.1463967](https://doi.org/10.1080/10408398.2018.1463967)

To link to this article: <https://doi.org/10.1080/10408398.2018.1463967>



© 2018 The Author(s). Published with license by Taylor & Francis Group, LLC
Lukas Schwingshackl, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann, and Heiner Boeing



[View supplementary material](#)



Published online: 11 May 2018.



[Submit your article to this journal](#)



Article views: 12435



[View related articles](#)



[View Crossmark data](#)



Citing articles: 33 [View citing articles](#)

REVIEW



Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: A systematic review and network meta-analysis

Lukas Schwingshackl^{a,b}, Anna Chaimani^{c,d,e}, Carolina Schwedhelm^{a,b}, Estefania Toledo^{f,g,h}, Marina Pünsch^a, Georg Hoffmannⁱ, and Heiner Boeing^{a,b}

^aDepartment of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Arthur-Scheunert-Allee 114–116, Nuthetal, Germany; ^bNutriAct-Competence Cluster Nutrition Research Berlin-Potsdam, Nuthetal, Germany; ^cParis Descartes University, Paris, France; ^dINSERM, UMR1153 Epidemiology and Statistics, Sorbonne Paris Cité Research Center (CRESS), METHODS Team, Paris, France; ^eCochrane France, Paris, France; ^fDepartment of Preventive Medicine and Public Health, Universidad de Navarra, Calle Irunlarrea 1, Pamplona, Spain; ^gCentro del Investigación Biomédica en Red de la Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Av. Monforte de Lemos 3–5, Madrid, Spain; ^hIdiSNA, Navarra Institute for Health Research, Calle Irunlarrea 3, Pamplona, Spain; ⁱDepartment of Nutritional Sciences, University of Vienna, Althanstraße 14, Vienna, Austria

ABSTRACT

Background: Pairwise meta-analyses have shown beneficial effects of individual dietary approaches on blood pressure but their comparative effects have not been established. **Objective:** Therefore we performed a systematic review of different dietary intervention trials and estimated the aggregate blood pressure effects through network meta-analysis including hypertensive and pre-hypertensive patients. **Design:** PubMed, Cochrane CENTRAL, and Google Scholar were searched until June 2017. The inclusion criteria were defined as follows: i) Randomized trial with a dietary approach; ii) hypertensive and pre-hypertensive adult patients; and iii) minimum intervention period of 12 weeks. In order to determine the pooled effect of each intervention relative to each of the other intervention for both diastolic and systolic blood pressure (SBP and DBP), random effects network meta-analysis was performed. **Results:** A total of 67 trials comparing 13 dietary approaches (DASH, low-fat, moderate-carbohydrate, high-protein, low-carbohydrate, Mediterranean, Palaeolithic, vegetarian, low-GI/GL, low-sodium, Nordic, Tibetan, and control) enrolling 17,230 participants were included. In the network meta-analysis, the DASH, Mediterranean, low-carbohydrate, Palaeolithic, high-protein, low-glycaemic index, low-sodium, and low-fat dietary approaches were significantly more effective in reducing SBP (–8.73 to –2.32 mmHg) and DBP (–4.85 to –1.27 mmHg) compared to a control diet. According to the SUCRA, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP). For most comparisons, the credibility of evidence was rated very low to moderate, with the exception for the DASH vs. the low-fat dietary approach for which the quality of evidence was rated high. **Conclusion:** The present network meta-analysis suggests that the DASH dietary approach might be the most effective dietary measure to reduce blood pressure among hypertensive and pre-hypertensive patients based on high quality evidence.

KEYWORDS

Dietary approaches; hypertension; network meta-analysis; blood pressure; evidence synthesis

1. Background

According to the World Health Organization, approximately 1 billion adults worldwide suffer from hypertension, and the prevalence is increasing (NCD-Risk-Factor-Collaboration 2017; World-Health-Organization 2012). Hypertension is the most important modifiable risk factor for premature cardiovascular disease worldwide. Nearly 50% of all ischemic heart diseases and stroke events are attributable to hypertension (Lawes, Vander Hoorn, and Rodgers 2008).

Dietary modifications play an important role in the prevention and management of hypertension. Evidence from a recent dose-response meta-analysis of prospective cohort studies indicates that optimal consumption of certain food groups (whole grains, fruit, nuts, legumes, dairy products, red and processed

meat and sugar sweetened beverages) is associated with as strong reduction in hypertension risk (Schwingshackl et al. 2017b). According to the most recent guidelines by the American Heart Association, hypertensive and pre-hypertensive patients should follow dietary modifications (alcohol moderation, sodium reduction, and emphasis on increased consumption of fresh fruits, vegetables, and low-fat dairy products) (Eckel et al. 2014; Smith et al. 2011). However, considering various guidelines, dietary recommendations for the prevention and management of hypertension are still not comprehensive (Eckel et al. 2014; Mancia et al. 2013b; Smith et al. 2011).

Pairwise meta-analyses comparing Dietary approaches to stop hypertension (DASH) (Saneii et al. 2014), combined

CONTACT Lukas Schwingshackl, PhD ✉ lukas.schwingshackl@dife.de 📧 Arthur-Scheunert-Allee 114–116, Nuthetal, 14558, Germany.

Financial support: No funding to declare.

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/bfsn.

Supplemental data for this article can be accessed on the [publisher's website](http://www.tandfonline.com/bfsn).

© 2018 Lukas Schwingshackl, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann, and Heiner Boeing. Published with license by Taylor & Francis Group, LLC. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

dietary approaches (Ndanuko et al. 2016), or lower sodium intake with usual care/control diets, indicate modest reductions in blood pressure (Aburto et al. 2013). Due to the high prevalence of hypertension, and the preventive effects of dietary factors, one of the most important questions that remain to be answered is which dietary approach offers the greatest benefits. For answering this question, a promising approach is network meta-analysis (Leucht et al. 2016; Mavridis et al. 2015; Salanti 2012). The methodology of network meta-analysis is an extension of the standard pairwise meta-analysis that enables a simultaneous comparison of multiple interventions forming a connected network while preserving the internal randomization of individual trials. This systematic review and network meta-analysis aims to establish a clinically meaningful hierarchy of different dietary approaches on blood pressure in hypertensive patients in a systematic review through the synthesis of available evidence from randomized trials.

2. Methods and design

The study protocol was registered in PROSPERO International Prospective Register of Systematic Reviews (<https://www.crd.york.ac.uk/prospero/>, identifier CRD42016049243) and published a priori (Schwingshackl et al. 2017a). The present systematic review adheres to the PRISMA guidelines and the corresponding extension for network meta-analyses (Chaimani et al. 2017; Hutton et al. 2015).

2.1. Search strategy

Literature searches were performed in PubMed, Cochrane Central Register of Controlled Trials (CENTRAL), and Google Scholar until June 2017, by applying no language restriction using a pre-defined search strategy (Supplemental Appendix 1).

Moreover, the reference lists from the identified articles were screened to search for additional relevant studies.

2.2. Eligibility criteria

As previously described the following inclusion criteria were established (Schwingshackl et al. 2017a):

- i) Randomized comparison/controlled design (parallel or cross-over) between different dietary approaches (Schwingshackl et al. 2018):
 - a. DASH diet: high intake of fruits & vegetables, low-fat dairy products, and whole grains, and low in sodium (Appel et al. 1997);
 - b. Mediterranean dietary pattern: high consumption of fruit, vegetables, olive oil, legumes, cereals, fish, and moderate intake of red wine during meals (Schwingshackl and Hoffmann 2014a, 2014c, 2015a, 2016; Schwingshackl et al. 2015b);
 - c. Low carbohydrate (LC) diet: <25% carbohydrates of total energy intake; high intake of animal and/ or plant protein; often high intake of fat (Schwingshackl and Hoffmann 2013a);
 - d. Palaeolithic Diet: lean meat, fish, fruit, leafy and cruciferous vegetables, root vegetables, eggs and nuts, while excluding dairy products, cereal grains, beans,

refined fats, sugar, candy, soft drinks, beer and extra addition of salt (Jonsson et al. 2009);

- e. Moderate-carbohydrate diet: 25–45% carbohydrates of total energy intake; 10–20% protein intake (Schwingshackl and Hoffmann 2013a);
- f. High protein (HP) diet: >20% protein intake of total energy intake; high intake of animal and/ or plant protein; <35% fat (Schwingshackl and Hoffmann 2013b);
- g. Nordic Diet: whole-grain products, abundant use of berries, fruit and vegetables, rapeseed oil, three fish meals per week, low-fat dairy products and avoidance of sugar-sweetened products (Uusitupa et al. 2013);
- h. Tibetan Diet: high-protein and vitamin-rich food, preferably cooked and warm food (von Haehling et al. 2013);
- i. Low fat (LF) diet: <30% fat of total energy intake; high intake of cereals & grains; 10–15% protein intake (Schwingshackl and Hoffmann 2013a, 2014b);
- j. Low glycaemic index/load (LGI/LGL) diet (Schwingshackl, Hobl, and Hoffmann 2015a; Schwingshackl and Hoffmann 2013c);
- k. Vegetarian/Vegan diet: no meat and fish/ no animal products (Haider et al. 2017)
 - l. Low-sodium diet (Whelton et al. 1998);
 - m. Control diet: usual diet.
- ii) Minimum intervention period of 12 weeks;
- iii) Participants with a mean age ≥ 18 years;
- iv) Hypertension or pre-hypertension (high normal blood pressure: 130–139 mmHg for systolic and/or 85–89 for diastolic blood pressure) defined according to the European Society of Hypertension and the European Society of Cardiology (Mancia et al. 2013a);
- v) The outcomes include systolic (SBP) and diastolic blood pressure (DBP) (mmHg).

The following studies were excluded:

- i) Randomized trials including pregnant women, children, and adolescents;
- ii) Intervention studies solely based on dietary supplements or single foods;
- iii) Intervention studies using dietary supplements as placebo;
- iv) Studies with an exercise/medication (Schwingshackl et al. 2013; Schwingshackl et al. 2014) co-intervention that was not applied in all the intervention/control groups;
- v) Interventions based on very low energy diets (i.e. <600 kcal/day);

2.3. Data extraction

After determination of the study selection, two reviewers extracted the following characteristics: name of first author, year of publication, country, sample size, study design, mean baseline Body Mass Index (BMI), mean baseline SBP and DBP, mean baseline age, % female, study duration, description of the different dietary arms, type of diet (energy restricted, ad libitum, iso-caloric), drop outs, % hypertension, % presence of comorbidities (obesity, hyperlipidaemia, type 2 diabetes,

metabolic syndrome, coronary heart disease), % antihypertensive medication, and % lipid lowering medication. Outcome data include: post-intervention values with corresponding standard deviations for SBP and DBP.

2.4. Risk of bias assessment

One author assessed the risk of bias of the included trials, by applying the Cochrane risk of bias tool (Higgins et al. 2011). The following items were assessed: random sequence generation and allocation concealment, blinding of participants and personnel, incomplete outcome data, and selective reporting.

Studies were classified as being at low risk of bias (if at least three out of a maximum of five items were rated as low risk; and maximum one item rated with a high risk of bias), high risk of bias (if at least two out of a maximum of five items were rated as high risk), and moderate/unclear risk (all other studies).

2.5. Dealing with missing data

We contacted authors to receive missing outcome data (1 author sent additional data). If the post-intervention values with the corresponding standard deviations were not available, the change scores with the corresponding standard deviations were used, according to the guidelines of the Cochrane Handbook (Higgins and Green 2011).

2.6 Data synthesis

2.6.1 Description of the available data

Direct comparisons between different dietary approaches were illustrated by using a network diagram (Chaimani et al. 2013). Moreover, the contribution matrix was used to identify the direct comparisons with greater influence in the network relative effects (Chaimani et al. 2013; Krahn, Binder, and Konig 2013).

2.6.2 Assessment of transitivity

To evaluate the assumption of transitivity we compared the distribution of the potential effect modifiers (age, study length, Body Mass Index (BMI)) across the available direct comparisons.

2.6.3 Statistical analysis

In order to quantify the pooled relative effect of each dietary approach against every other dietary approach in terms of the post-intervention values (or changes from baseline values) of the different dietary measures we performed a random effects network meta-analysis for SBP and DBP. We performed network meta-analysis to synthesize all the available evidence. Compared to the standard pairwise meta-analysis model, network meta-analysis methods are extensions that enable a simultaneous comparison of multiple interventions forming a connected network while preserving the internal randomization of individual trials.

Random effects network meta-analysis for SBP and DBP were performed to estimate all possible pairwise relative effects and obtain a relative ranking of the different dietary approaches. The summary mean differences with their 95% CI were presented in a League table. We estimated the relative ranking of the different dietary approaches for SBP and DBP (Salanti, Ades, and Ioannidis 2011) with using the distribution of the ranking probabilities and the surface under the cumulative ranking curves (SUCRA). For SBP and DBP, we assumed a common network-specific heterogeneity parameter and estimated the predictive intervals to assess how much this heterogeneity affects the relative effects with respect to the additional uncertainty anticipated in future studies (Riley, Higgins, and Deeks 2011). Statistical analyses were performed by using Stata 14.0 (StataCorp. 2015) (*network* package (White 2015)) and produced presentation tools with the *network graphs* package (Chaimani and Salanti 2015).

2.6.4 Assessment of inconsistency

To evaluate the presence of statistical inconsistency, the loop-specific approach (Bucher et al. 1997) (detection of loops of evidence that might present important inconsistency), and the side-splitting approach (Dias et al. 2010) (detect comparisons for which direct estimates disagree with indirect evidence from the entire network) were applied. Global methods jointly investigate the presence of inconsistency from all possible sources in the entire network simultaneously. For this purpose, we used the design-by-treatment interaction model (Higgins et al. 2012; Jackson et al. 2014).

2.6.5 Subgroup and sensitivity analyses

Subgroup analyses according to hypertension status, study length (≥ 12 months vs. < 12 months), and sample size (≥ 100 vs. < 100) were performed for SBP and DBP. We also conducted a sensitivity analysis by analysing only studies considered being at low risk of bias. Additionally, we ran meta-regressions by participant and study characteristics for SBP and DBP: age, study length, and mean differences in weight change (by comparing study arms within each trial).

2.6.6 Small study effects and publication bias

The presence of small-study effects for each SBP and DBP was evaluated by drawing comparison-adjusted funnel plot that accounts for the fact that different studies compare different sets of interventions (Chaimani et al. 2013). Funnel plots included all comparisons of dietary approaches compared to a low-fat or control diet.

2.7 Credibility of the evidence

To assess the credibility of evidence from the network meta-analysis, we used the GRADE system extended for network meta-analysis following the approach suggested by Salanti et al. (Supplemental Appendix 2) (Salanti et al. 2014).

3. Results

Of a total of 2,949 records identified in the literature search, 115 full text articles were assessed in detail (**Supplemental Figure 1**). Of these, 48 were excluded, for the exclusion criteria summarized in **Supplemental Table 1**, **Supplemental References**.

Overall 67 trials met the eligibility criteria and were included in the meta-analysis (**Table 1**). 17,230 pre-hypertensive or hypertensive patients were enrolled, and the trials were published between 1981 and 2016. Eighteen trials were conducted in North America, 29 trials in Europe, 4 trials in Asia, and 15 trials in Australia and New Zealand, and one study in South America. Study length ranged between 3 and 48 months; the patients' mean age was between 23.6 and 71 years, and their BMI between 23.6 and 45.4 kg/m². The general and specific study characteristics are summarized in **Supplemental Table 2–3**.

Twenty-three trials were judged to be low risk of bias, five trials to be high risk of bias, and 39 trials were classified as moderate/unclear risk of bias studies. With regard to the specific items of the risk of bias assessment tool by the Cochrane Collaboration, 46% of the included studies indicate a low risk of bias for random-sequence generation, 21% for allocation concealment, 0% for blinding, 60% for incomplete data outcome, and 96% for selective reporting (**Supplemental Figure 2**).

The definition of the different intervention diets was heterogeneous for the intensity and type of dietary approach (i.e. group meeting, dietary counselling) and for the prescribed diets (ad-libitum, iso-caloric, hypocaloric). Across the included low-fat dietary approaches the overall amount of fat intake varied between ~10–15% of total energy intake. We thus had to harmonize the single trials and formed classes of dietary approaches (Schwingshackl et al. 2017a).

Figure 1 shows the network diagram of direct comparison for SBP with the number of studies reflected by the size of the edges, and the number of patients reflected by the size of the nodes. The comparisons with the largest amount of trials include: HP vs. LF (Brinkworth et al. 2004a; Brinkworth et al. 2004b; Campbell and Meckling 2012; Dalle Grave et al. 2013; Dansinger et al. 2005; Delbridge et al. 2009; Kim et al. 2014;




Figure 1. Network diagram for diastolic blood pressure: The size of the nodes is proportional to the total number of participants allocated to each dietary approach and the thickness of the lines proportional to the number of studies evaluating each direct comparison.

Krebs et al. 2012; Luger et al. 2013; Tang et al. 2013; Watson et al. 2016; Wycherley et al. 2010) ($n = 12$), low-sodium vs. control (Alli et al. 1992; Chalmers et al. 1986; Costa et al. 1981; Dodson et al. 1989; Erwtaman et al. 1984; Jula and Karanko 1994; Makela et al. 2008; Silman et al. 1983; Whelton 1997; Whelton et al. 1998) ($n = 10$), LF vs. control (Anderssen et al. 1995; Andrews et al. 2011; Ard et al. 2016; Coppell et al. 2010; Gordon, Scott, and Levine 1997; Heilbronn, Noakes, and Clifton 1999; Rock et al. 2014; Uusitupa et al. 1993; Watkins et al. 2003) ($n = 9$), and LC vs. LF dietary approaches (Daly et al. 2006; Guldbrand et al. 2012; Iqbal et al. 2010; Liu et al. 2013; Stern et al. 2004; Tay et al. 2008; Tay et al. 2015; Veum et al. 2017) ($n = 8$). **Table 2** shows the percentage of statistical contribution coming from direct and indirect comparisons for each dietary approach compared to each other. It was shown that most of the contribution to the study effects came from indirect comparisons. Direct comparisons dominated the comparisons of DASH, Vegetarian, Mediterranean, HP, moderate-carbohydrate, LC, LGI/LGL, Palaeolithic, Tibetan, and a control diet with a LF diet for both outcomes. In general, there are some important differences in the examined effect modifiers across comparisons for BMI and study duration. For some comparisons, though, we did not have enough studies and we could not test transitivity appropriately (**Supplemental Figure 3–5**).

Table 3 summarizes the pooled estimates of SBP and DBP for the comparison of every dietary approach using both direct and indirect evidence. The DASH, Mediterranean, LC, Palaeolithic, HP, LGI/LGL, low-sodium, and LF dietary approaches were more effective in reducing SBP and DBP compared to a control diet. The DASH diet was more effective in reducing SBP compared to a LF (mean difference (MD): -5.05 mmHg, 95% -7.08 , -3.03), Mediterranean (MD: -3.31 mmHg, 95% -6.20 , -0.42), HP (MD: -4.78 mmHg, 95% -7.72 , -1.85), moderate-carbohydrate (MD: -5.57 mmHg, 95% -8.75 , -2.39), LGI/LGL (MD: -4.92 mmHg, 95% -8.55 , -1.30), Nordic (MD: -4.74 mmHg, 95% -9.11 , -0.37), and low-sodium dietary approaches (MD: -4.42 mmHg, 95% -7.12 , -1.72), and more effective in reducing DBP compared to a LF diet (MD: -3.10 mmHg, 95% -4.52 , -1.68), moderate-carbohydrate (MD: -3.05 mmHg, 95% -5.37 , -0.74), Nordic

Table 1. References of the 67 trials included in the present network meta-analysis.

Alli et al. 1992; Anderssen et al. 1995; Andrews et al. 2011; Appel et al. 2003; Ard et al. 2016; Azadbakht et al. 2005; Blumenthal et al. 2010; Brehm et al. 2009; Brinkworth et al. 2004a; Brinkworth et al. 2004b; Campbell and Meckling 2012; Chalmers et al. 1986; Clifton et al. 2004; Coppell et al. 2010; Costa et al. 1981; Dalle Grave et al. 2013; Daly et al. 2006; Dansinger et al. 2005; Delbridge et al. 2009; Deluis et al. 2010; Dodson et al. 1989; Edwards et al. 2011; Erwtaman et al. 1984; Esposito et al. 2009; Esposito et al. 2004; Frisch et al. 2009; Gordon, Scott, and Levine 1997; Guldbrand et al. 2012; Heilbronn, Noakes, and Clifton 1999; Iqbal et al. 2010; Itsiopoulos et al. 2011; Jonsson et al. 2009; Jula and Karanko 1994; Kim et al. 2014; Kirpizidis, Stavratsi, and Geleris 2005; Klemsdal et al. 2010; Krebs et al. 2012; Lean et al. 1997; Lima et al. 2013; Liu et al. 2013; Luger et al. 2013; Ma et al. 2008; Makela et al. 2008; Mellberg et al. 2014; Nicholson et al. 1999; Nowson et al. 2005; Philippou et al. 2009; Poulsen et al. 2015; Rock et al. 2014; Shai et al. 2008; Silman et al. 1983; Stern et al. 2004; Tang et al. 2013; Tay et al. 2008; Tay et al. 2015; Toledo et al. 2013; Uusitupa et al. 2013; Uusitupa et al. 1993; Veum et al. 2017; von Haehling et al. 2013; Walker et al. 1995; Watkins et al. 2003; Watson et al. 2016; Westman et al. 2008; Whelton 1997; Whelton et al. 1998; Wycherley et al. 2010

Table 2. Percentage contribution of each direct estimate derived from direct (blue) and indirect (red) comparisons (the colour corresponds to the percentage of contribution). The values above the diet classes correspond to the percentage contribution of direct and indirect comparisons between the row and columns for DBP (eg, the percentage contribution of direct comparisons for DBP between Mediterranean and Low-Carb diet is 33%, and 65% for the indirect comparisons). The values below the diet classes correspond to the percentage contribution of direct and indirect comparisons between the column and the rows for SBP (eg, the percentage contribution of direct comparisons for SBP between Mediterranean and Low-Carb diet is 34%, and 66% for the indirect comparisons). DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.

DBP																				
DASH	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	59%	41%	60%	40%
0%	100%	Mediterranean	35%	65%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	87%	13%	0%	100%
0%	100%	34%	66%	Low-Carb	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	72%	28%	0%	100%
0%	100%	0%	100%	0%	100%	0%	100%	Palaeolithic	0%	100%	0%	100%	0%	100%	0%	100%	100%	0%	0%	100%
0%	100%	0%	100%	0%	100%	High-Protein	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	100%	0%	0%	100%
0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	Vegetarian	0%	100%	0%	100%	0%	100%	100%	0%	0%	100%
0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	Moderate-Carb	0%	100%	0%	100%	79%	21%	27%	73%
0%	100%	0%	100%	16%	84%	0%	100%	0%	100%	0%	100%	Low GI/GL	0%	100%	0%	100%	90%	10%	0%	100%
0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	Nordic	0%	100%	0%	100%	100%	0%	0%
0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	0%	Tibetan	0%	100%	100%	0%	0%	100%
0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	100%	0%	0%	0%	100%	Low-Sodium	0%	100%	100%	0%
62%	38%	87%	13%	73%	27%	100%	0%	100%	0%	100%	81%	19%	90%	10%	0%	100%	Low-Fat	74%	26%	26%
57%	43%	0%	100%	0%	100%	0%	100%	0%	100%	0%	27%	73%	0%	100%	0%	100%	73%	27%	Control	Control
SBP																				

Table 3. League table: The values above the diet classes correspond to the difference in mean (95% CI) in DBP (mmHg) between the row and columns (e.g. the mean difference in average DBP between DASH and Low-Fat diets diet is -3.10 mmHg). The value below the diet classes correspond to the difference in mean in SBP (mmHg) between the column and the row (e.g. the mean difference in average SBP between DASH and Low-Fat diet is -5.05 mmHg). DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.

		DBP (mmHg)													
DASH		-1.74	0.48	-1.91	-5.90	-3.05	-1.60	-3.22	-3.20	-1.91	-3.10	-4.37			
-3.31	Mediterranean	(-3.92,0.43)	(-3.72,4.68)	(-3.92,0.10)	(-19.60,7.80)	(-5.37,-0.74)	(-4.05,0.85)	(-6.14,-0.31)	(-6.32,-0.08)	(-3.70,-0.12)	(-4.52,-1.68)	(-5.78,-2.97)			
-2.86		-0.18	2.04	-0.35	4.34	-1.50	-0.04	-1.66	-1.64	-0.35	-1.54	-2.81			
		(-2.11,1.75)	(-2.16,6.24)	(-2.35,1.65)	(-18.04,9.36)	(-3.85,0.86)	(-2.47,2.39)	(-4.80,1.47)	(-4.75,1.47)	(-2.47,1.76)	(-2.95,-0.13)	(-4.63,-1.00)			
	Low-Carb	2.22	2.22	-0.17	-4.16	-1.31	0.14	-1.48	-1.46	-0.17	-1.36	-2.63			
(-5.96,0.25)		(-2.35,2.01)	(-2.07,6.51)	(-2.35,2.01)	(-17.89,9.57)	(-3.82,1.19)	(-2.34,2.63)	(-4.73,1.77)	(-4.69,1.77)	(-2.45,2.11)	(-3.01,0.30)	(-4.64,-0.62)			
1.35		4.21		-2.39	-6.38	-3.53	-2.08	-3.70	-3.68	-2.39	-3.58	-4.85			
(-6.31,9.01)	Palaeolithic	(-3.55,11.97)		(-6.59,1.82)	(-20.57,7.81)	(-7.92,0.85)	(-6.51,2.35)	(-8.55,1.14)	(-8.51,1.16)	(-6.66,1.88)	(-7.53,0.38)	(-8.97,-0.73)			
-4.78		-1.92	-13.83	High-Protein	-3.99	-1.15	0.31	-1.31	-1.29	-0.00	-1.19	-2.46			
(-7.72,-1.85)		(-5.11,1.26)	(-13.83,1.56)		(-17.69,9.71)	(-3.51,1.22)	(-2.14,2.76)	(-4.45,1.82)	(-4.41,1.83)	(-2.14,2.14)	(-2.61,0.23)	(-4.28,-0.64)			
-0.65		2.20	-2.00	4.13	Vegetarian	2.84	4.30	2.68	2.70	3.99	2.80	1.53			
(-17.15,15.84)		(-14.34,18.75)	(-19.97,15.96)	(-12.38,20.64)		(-10.92,16.60)	(-9.47,18.07)	(-11.24,16.59)	(-11.21,16.61)	(-9.74,17.71)	(-10.83,16.43)	(-12.15,15.20)			
-5.57		-2.71	-6.92	-0.79	-4.91	Moderate-Carb	1.46	-0.17	-0.14	1.14	-0.04	-1.32			
(-8.75,-2.39)		(-6.19,0.76)	(-14.74,0.90)	(-4.11,2.53)	(-21.48,11.66)		(-1.29,4.20)	(-3.48,3.14)	(-3.50,3.21)	(-1.24,3.52)	(-1.93,1.84)	(-3.42,0.78)			
-4.92		-2.07	-6.27	-0.14	-4.27		Low GI/GL	-1.62	-1.60	-0.31	-1.50	-2.77			
(-8.55,-1.30)		(-5.74,1.61)	(-14.26,1.71)	(-3.83,3.55)	(-20.92,12.38)	0.65		(-5.06,1.81)	(-5.02,1.82)	(-2.88,2.25)	(-3.49,0.49)	(-5.06,-0.48)			
-4.74		-1.88	-6.09	0.04	-4.08		0.19	Nordic	0.02	1.31	0.12	-1.15			
(-9.11,-0.37)		(-6.70,2.94)	(-14.59,2.41)	(-4.66,4.75)	(-20.99,12.82)	0.83			(-3.91,3.96)	(-1.49,4.11)	(-2.67,2.92)	(-3.70,1.41)			
-3.95		-0.64	-5.30	0.83	-3.30		0.97		(-3.91,3.96)	1.29	0.10	-1.17			
(-8.48,0.57)		(-5.79,3.59)	(-13.73,3.12)	(-3.75,5.40)	(-20.17,13.57)	1.61		0.78	Tibetan	1.29	0.10	(-3.70,1.41)			
-4.42		-1.57	-5.77	0.36	-3.77		0.50	(-5.05,6.61)		(-1.92,4.49)	(-2.68,2.88)	(-4.17,1.83)			
(-7.12,-1.72)		(-4.92,1.79)	(-13.55,2.00)	(-2.86,3.58)	(-20.32,12.78)	1.15	0.31	0.31	-0.47	Low-Sodium	(-2.79,0.41)	(-3.59,-1.32)			
-5.05		-2.20	-6.40	-0.27	-4.40		(-3.38,4.38)	(-3.93,4.56)	(-5.19,4.25)	-0.63	Low-Fat	(-3.59,-1.32)			
(-7.08,-3.03)		(-4.56,0.17)	(-13.80,0.99)	(-2.40,1.86)	(-20.77,11.97)	0.51	-0.13	(-4.51,3.88)	(-5.15,2.95)	(-3.06,1.80)		(-2.41,-0.14)			
-7.38		-4.52	-8.73	-2.60	-6.72		-2.46	-2.64	-3.42	-2.96	-2.32	(-2.41,-0.14)			
(-9.43,-5.33)		(-7.41,-1.64)	(-16.31,-1.15)	(-5.29,0.10)	(-23.18,9.73)		(-4.69,1.07)	(-6.50,1.22)	(-7.80,0.95)	(-4.74,-1.17)	(-3.98,-0.67)	Control			

(MD: -3.22 mmHg, 95% -6.14 , -0.31), Tibetan (MD: -3.20 mmHg, 95% -6.32 , -0.08) and low-sodium diet (MD: -1.91 mmHg, 95% -3.70 , -0.12). The Mediterranean dietary approach was more effective in reducing DBP compared to a LF diet (MD: -1.54 mmHg, 95% -2.95 , -0.13). According to the SUCRAs, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP) (Supplemental Table 4–5).

The rankograms showed uncertainty in ranking (similar distribution of rank probabilities across many possible ranks indicates uncertain ranking for that dietary approach) for the HP, LC, LGI/LGL, moderate-carbohydrate, Mediterranean, low-sodium, Nordic, and Tibetan dietary approach compared to the other dietary approaches (Supplemental Figure 6–7).

The side-splitting approach suggested no significant inconsistency for SBP and DBP (Supplemental Table 6–7). The loop-specific approach identified two-loops (control – LF – moderate-carbohydrate; and LF – LC – Mediterranean) for SBP presenting statistical inconsistency, and one-loop for DBP (LF – LC – LGI/GL) (Supplemental Figure 8–9). The design-by-treatment model did not suggest the presence of statistical inconsistency for SBP ($p = 0.65$), and DBP ($p = 0.83$). The important inconsistency in the loop specific approach might be explained by several differences across LF dietary approaches (hypocaloric if compared to a control diet; often iso-caloric if compared to other dietary approaches), differences in total fat to carbohydrate intake ratio, and fatty acids composition among moderate carbohydrate approaches, and LC dietary approaches (larger weight loss compared to other interventions).

3.1 Subgroup and sensitivity analysis

In sensitivity analysis including only studies with a low risk of bias, the results of the primary analysis for the best ranked dietary approach, i.e. the DASH diet could be confirmed (Table 4). No significant effects for all other dietary approaches were observed in the low risk of bias sensitivity analysis.

The a priori planned subgroup analysis for age was not possible to conduct since only 8 trials included participants >60 years of age. In the subgroup analyses comparing longer-term (≥ 12 months) vs. shorter-term trials (<12 months), most of the results of the primary analysis were not confirmed among longer-term studies. Moreover, it seems that trials with a smaller sample size (<100) yielded more significant results compared to studies with a larger sample size (≥ 100) (Supplemental Table 8–15).

Due to the low number of studies presenting results for hypertensive patients ($n = 12$), it was not possible to conduct sensitivity analysis.

In univariate meta-regression analysis we could show that mean reduction in SBP and DBP was larger in short-term trial, and among trials including younger participants. Moreover, we could show that larger differences in body weight change between study arms within trials were associated with a stronger impact on SBP and DBP, compared to smaller differences

(Supplemental Figure 10–15), thereby showing that weight loss is a major contributing factor for blood pressure control.

Small study effects

The comparison-adjusted funnel plots for SBP and DBP appeared asymmetric suggesting that small studies tend to favour the dietary approaches when compared to control dietary approaches, and slightly asymmetric for low fat diets (Supplemental Figure 16–19).

3.2 Credibility of the evidence

For most comparisons in the network meta-analysis, the credibility of evidence was rated very low to moderate with the exception for the DASH vs. LF dietary approach the quality of evidence was rated high (Supplemental Figure 20, Supplemental Appendix 2).

4. Discussion

In this meta-analysis of 67 trials including 17230 hypertensive and pre-hypertensive patients we compared the effects of 13 different dietary approaches (DASH, Control, Low-Fat, Moderate-carbohydrate, High-protein, Low-carbohydrate, Mediterranean, Palaeolithic, Vegetarian, Low-GI/GL, Low-sodium, Nordic, and Tibetan) using network meta-analysis. According to the SUCRAs, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP). The DASH, Mediterranean, low-carbohydrate, Palaeolithic, high-protein, low-glycaemic index, low-sodium, and low-fat dietary approach were significantly more effective in reducing SBP (-8.73 to -2.32 mmHg) and DBP (-4.85 to -1.27 mmHg) compared to a control diet. For most comparisons, the credibility of evidence was rated very low to moderate, with the exception for the DASH vs. the low-fat dietary approach the quality of evidence was rated high.

Compared to a LF or a control diet, the DASH dietary approach reduced SBP by approximately 5–7 mmHg, and DBP by 3–4 mmHg. With respect to other therapeutic options, a recent meta-analysis investigating the effect of different exercise modalities resulted in some less pronounced effect of aerobic exercise on blood pressure (Cornelissen and Smart 2013). Moreover, the blood-pressure lowering effect of the DASH dietary pattern in our meta-analysis was comparable to drug monotherapy (Elmer et al. 2006).

The importance of blood pressure reduction, as shown for several dietary approaches is strengthened by a large meta-analysis of epidemiological studies which have shown that a decrease of approximately 10 mmHg reduction in SBP was inversely associated with risk of cardiovascular disease events by 20%, coronary heart disease by 17%, stroke by 27%, heart failure by 28% and all-cause mortality by 13% (Ettehad et al. 2016), whereas a 5 mmHg decrease in DBP reduces the risk of ischemic heart disease by 20%, and the risk of stroke by 32% (Law, Wald, and Morris 2003). Another meta-analysis stressed the fact that even a small decline in SBP of about 2 mmHg will be accompanied by a 10% lower risk of death due to stroke or 7% due to ischemic heart disease (Lewington et al. 2002). Albeit

Table 4. League table (Low risk of bias sensitivity analysis): The values above the diet classes correspond to the difference in mean (95% CI) in DBP (mmHg) between the row and columns (eg, the mean difference in average DBP between DASH and Low-Fat diets is -3.18 mmHg). The value below the diet classes correspond to the difference in mean in SBP (mmHg) between the column and the row (eg, the mean difference in average SBP between DASH and Low-Fat diets is -5.83 mmHg); DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.

	DBP (mmHg)										SBP (mmHg)	
DASH												
-4.15	-1.91 (-4.00, 0.18)	-2.59 (-5.05, -0.13)	0.82 (-5.55, 7.19)	-1.83 (-4.55, 0.88)	-4.01 (-7.76, -0.27)	-2.80 (-5.98, 0.38)	-1.09 (-4.90, 2.72)	-3.42 (-5.83, -1.02)	-3.18 (-4.89, -1.47)	-3.79 (-5.72, -1.86)		
-4.24	-0.08 (-3.18, 3.01)	-0.68 (-2.62, 1.26)	2.73 (-3.55, 9.00)	0.07 (-2.41, 2.56)	-2.10 (-5.79, 1.58)	-0.89 (-3.84, 2.06)	0.82 (-3.06, 4.69)	-1.52 (-4.02, 0.99)	-1.27 (-2.59, 0.05)	-1.88 (-3.93, 0.16)		
3.17	7.32 (-6.81, 21.46)	7.41 (-6.86, 21.67)	3.41 (-2.99, 9.81)	0.76 (-2.04, 3.55)	-1.42 (-5.32, 2.47)	-0.21 (-3.24, 2.83)	1.50 (-2.58, 5.57)	-0.83 (-3.64, 1.97)	-0.59 (-2.42, 1.24)	-1.20 (-3.61, 1.21)		
-7.12	-2.96 (-7.36, 1.43)	-2.88 (-7.69, 1.93)	Palaeolithic (-10.29, 24.78, 4.20)	-2.65 (-9.15, 3.84)	-4.83 (-11.86, 2.20)	-3.62 (-10.30, 3.07)	-1.91 (-9.05, 5.22)	-4.24 (-10.74, 2.25)	-4.00 (-10.13, 2.13)	-4.61 (-10.94, 1.72)		
-6.96	-2.81 (-8.32, 2.71)	-2.73 (-8.58, 3.13)	-10.13 (-25.00, 4.74)	0.16 (-6.23, 6.54)	-2.18 (-6.23, 1.87)	-0.96 (-4.39, 2.46)	0.74 (-3.48, 4.97)	-1.59 (-4.61, 1.43)	-1.35 (-3.49, 0.80)	-1.96 (-4.61, 0.70)		
-2.96	1.19 (-3.72, 6.10)	1.27 (-3.69, 6.24)	-6.13 (-20.81, 8.54)	4.15 (-1.76, 10.06)	4.00 0 (-2.79, 10.79)	1.21 (-3.13, 5.56)	2.92 (-1.87, 7.71)	0.59 (-3.18, 4.36)	0.83 (-2.61, 4.27)	0.22 (-3.27, 3.71)		
-3.18	0.97 (-5.47, 7.41)	1.05 (-5.67, 7.78)	-6.35 (-21.59, 8.89)	3.94 (-3.26, 11.13)	3.78 (-3.72, 11.28)	Low GI/GL (-0.22, -7.78, 7.34)	1.71 (-2.80, 6.22)	-0.63 (-4.03, 2.78)	-0.38 (-3.04, 2.27)	-0.99 (-4.08, 2.09)		
-6.15	-2.00 (-6.04, 2.04)	-1.92 (-6.41, 2.57)	-9.32 (-23.71, 5.06)	0.96 (-4.20, 6.13)	0.81 (-4.77, 6.39)	-3.19 (-8.84, 2.46)	Nordic (-2.97, -9.04, 3.09)	-2.33 (-5.92, 1.25)	-2.09 (-5.73, 1.55)	-2.70 (-5.99, 0.59)		
-5.83	-1.68 (-3.81, 0.46)	-1.59 (-4.49, 1.30)	-9.00 (-22.97, 4.97)	1.29 (-2.56, 5.14)	1.13 (-3.97, 6.23)	-2.87 (-7.35, 1.61)	-2.65 (-8.73, 3.44)	0.32 (-3.12, 3.77)	0.24 (-1.88, 2.37)	-0.37 (-1.80, 1.07)		
-6.88	-2.73 (-5.97, 0.51)	-2.65 (-6.43, 1.14)	-10.05 (-24.24, 4.13)	0.24 (-4.33, 4.80)	0.08 (-4.95, 5.11)	-3.92 (-9.03, 1.20)	-3.70 (-9.26, 1.86)	-0.73 (-3.14, 1.68)	-1.05 (-3.51, 1.41)	-0.61 (-2.18, 0.96)		
									Low-Fat	Control		

the overall net effect was modest for several dietary approaches, it is important to note that the control/low-fat groups also experienced some benefit and the results were incremental reductions experienced by those groups adopting dietary interventions.

Although this is the first network meta-analysis to assess the comparative effects of different dietary approaches, several previous pairwise meta-analyses have been published, either including only one dietary approach or mixing normotensive and hypertensive patients. A recent standard pairwise meta-analysis of 24 randomized trials including 23858 normotensive and hypertensive participants showed that the DASH, the Mediterranean, a low-sodium and a low-calorie diet were all effective in reducing blood pressure, whereas the Mediterranean diet did not significantly decrease SBP compared to control diets (Gay et al. 2016). Similar to our findings, previous meta-analyses reported a strong reduction in SBP and DBP by the DASH diet (Ndanuko et al. 2016; Saneai et al. 2014; Siervo et al. 2015).

The DASH dietary pattern is based on several food groups like whole grains, fruits, vegetables, dairy products (mainly low-fat), and nuts and lower intakes of red and processed meat as well as and sweets (Schwingshackl and Hoffmann 2015b). Optimal consumption of risk-decreasing foods (whole grains, fruits, nuts, legumes and dairy products) resulted in a 41% reduction of hypertension compared to non-consumption of these foods (Schwingshackl et al. 2017b). One potential explanation for the superiority of the DASH diet compared to the Mediterranean diet is the recommendation to increase dairy products and at the same time to lower sodium intake (Sacks et al. 2001) in the former regime, whereas the latter discourages higher intakes of dairy products and does not consider sodium in the diet (Trichopoulou et al. 2003). The original DASH diet was tested independent of sodium intake (Appel et al. 1997). However, 6 out of the included 7 trials investigating the DASH approach in the present NMA tested the effects of the refined DASH diet implementing sodium restriction, whereas one trial did not provide the corresponding information (Edwards et al. 2011).

Anti-hypertensive effects of DASH dietary components are biologically plausible, given the high intake of constituents like phytochemicals and nutrients such as magnesium, potassium, calcium, lactotripeptides, selenium, zinc, antioxidants, vitamins, unsaturated fatty acid, and fibre which have been shown to lower blood pressure (Alonso et al. 2006; Appel et al. 2006; Fekete, Givens, and Lovegrove 2015; Han et al. 2017; Houston 2011; Schwingshackl, Strasser, and Hoffmann 2011; Whelton et al. 2005).

Whereas the evidence for an antihypertensive effect of the DASH diet is highly consistent, inconsistent results were reported for the Mediterranean Diet. One recent meta-analysis of six trials including more than 7,000 participants found insufficient evidence to suggest that the Mediterranean Diet decreased blood pressure (Nissensohn et al. 2016). In contrast, another recent meta-analysis showed that the Mediterranean dietary approach was effective in reducing SBP and DBP (Ndanuko et al. 2016).

Across all dietary approaches, only the low-sodium diet was rated with a high quality of meta-evidence. The blood pressure lowering effect observed in the present meta-analysis is consistent with several pairwise meta-analyses of randomized trials

(Aburto et al. 2013). In line with our and previous findings the European Society of Hypertension and Cardiology rated the recommendation for the prevention and management of hypertension of a maximum daily salt intake of 5–6 gram with the highest level of evidence (Mancia et al. 2013b).

Previous meta-analyses reported conflicting results regarding the blood pressure lowering effects of LC and HP diets (Bueno et al. 2013; Mansoor et al. 2016; Santesso et al. 2012; Schwingshackl and Hoffmann 2013b; Wycherley et al. 2012). Although in the present network meta-analysis the LC and the HP diets were more effective in reducing blood pressure compared to control diets, the effects compared to LF diets were not statistically significant.

The results based on the Palaeolithic, Vegetarian, Nordic, and Tibetan dietary approach should be interpreted with caution, since only a very low number of trials (≤ 2) were available. Nevertheless, our findings for the Palaeolithic diet are in line with a previous pairwise meta-analysis showing a reduction in SBP (Manheimer et al. 2015). Blood pressure reductions of the Vegetarian diet and the Nordic diet as reported by previous meta-analyses could not be confirmed (Ndanuko et al. 2016; Yokoyama et al. 2014) in pre-hypertensive and hypertensive individuals. Similar to our findings, a recent meta-analysis comparing LGI/LGL vs. high-GI diets showed evidence of a blood pressure reducing effect of low GI/GL diets (Evans et al. 2017).

Beside diet and physical activity, but very likely mediated via these measures, weight loss is the third big lifestyle pillar in the prevention and management of hypertension (Mancia et al. 2013a). In a comprehensive dose-response meta-analysis, each kilogram of weight loss showed a reduction of approximately 1 mmHg of diastolic and systolic blood pressure, respectively (Neter et al. 2003). In line with these observations, the meta-regression analysis showed that larger differences in body weight changes between arms within trials were associated with a stronger impact on SBP and DBP, compared to smaller differences.

4.1. Strengths & limitations

The present systematic review has several strengths including the application of network meta-analyses methodology, which simultaneously combined direct and indirect evidence, the stringent inclusion and exclusion criteria, the high number of included dietary approaches, participants and trials, the comprehensive literature search, the assessment of the risk of bias, subgroup and sensitivity analyses, the a priori published systematic review protocol, the quality of meta-evidence assessment, and the overall low observed heterogeneity.

Several limitations should be considered when interpreting the results. First, all analyses were based on the original intended randomized design, not by adherence to the actual dietary approach and/or macronutrient composition and caloric intake consumed. This means that although participants were randomized to various dietary approaches, details on their actual adherence to the dietary program were not accounted for in the analyses. Second, 44 of 67 trials were rated as unclear or high risk of bias mostly due to lack of allocation concealment and blinding. The risk of bias sensitivity analysis was based on

23 trials, and only the primary results for the DASH dietary pattern could be confirmed. Third, was the substantial clinical heterogeneity for the included populations in terms of age, sex, study methods, study duration, sample size, and differences in food intake included across the various dietary pattern. Blood pressure reduction was lower in long-term studies, suggesting that participant's adherence decline over time. Moreover, larger size studies (sample size: ≥ 100) had a lower effect on blood pressure reduction compared to studies with a smaller sample size (< 100). Fourth, due to the low number of studies including hypertensive patients, we were not able to run a sensitivity network meta-analysis. Finally, several nodes were not well connected, due to the low number of trials and this should be taken into account when interpreting the results of the present network meta-analysis.

5. Conclusion

Our network meta-analysis suggest that the DASH dietary approach might be the most effective dietary approach to improve blood pressure in pre-hypertensive and hypertensive patients based on high quality evidence. The findings of the present network meta-analysis have important clinical and public health implications, suggesting that dietary modifications characterized by high consumption of fruit and vegetables, whole grains, legumes, nuts and seeds, and dairy products and low consumption of red and processed meat, sugar sweetened beverages and sodium are an effective method for controlling blood pressure within pre-hypertensive and hypertensive populations.

Competing interests

Authors declare to have no competing of interest.

Acknowledgments

We thank Thomas Wolever, PhD for sending us additional data for the corresponding meta-analysis.

Authors' contributions

LS, AC, HB, GH contributed to the conception and design of the systematic review and meta-analysis. LS, MP, HB, were involved in the acquisition and analysis of the data. LS, CS, HB, interpreted the results. LS, AC, GH, CS, ET, HB, drafted this manuscript. All authors provided critical revisions of the protocol and approved submission of the final manuscript.

ORCID

Lukas Schwingshackl  <http://orcid.org/0000-0003-3407-7594>

References

Aburto, N. J., A. Ziolkovska, L. Hooper, P. Elliott, F. P. Cappuccio, and J. J. Meerpohl. 2013. Effect of lower sodium intake on health: systematic review and meta-analyses. *Bmj (Clinical Research Ed.)* 346:f1326.

Alli, C., F. Avanzini, G. Bettelli, M. Bonati, F. Colombo, R. Corso, M. Di Tullio, M. G. Gentile, L. Sangalli, E. Taioli, et al. 1992. Feasibility of a long-term low-sodium diet in mild hypertension. *Journal of Human Hypertension* 6:281–86.

Alonso, A., J. J. Beunza, M. Bes-Rastrollo, R. M. Pajares, and M. A. Martinez-Gonzalez. 2006. Vegetable protein and fiber from cereal are inversely associated with the risk of hypertension in a Spanish cohort. *Archives of Medical Research* 37:778–86. doi:10.1016/j.arcmed.2006.01.007.

Anderssen, S., I. Holme, P. Urdal, and I. Hjermann. 1995. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press* 4: 343–49.

Andrews, R. C., A. R. Cooper, A. A. Montgomery, A. J. Norcross, T. J. Peters, D. J. Sharp, N. Jackson, K. Fitzsimons, J. Bright, K. Coulman, et al. 2011. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. *Lancet* 378:129–39. doi:10.1016/S0140-6736(11)60442-X.

Appel, L. J., M. W. Brands, S. R. Daniels, N. Karanja, P. J. Elmer, and F. M. Sacks. 2006. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension* 47:296–308. doi:10.1161/01.HYP.0000202568.01167.B6.

Appel, L. J., C. M. Champagne, D. W. Harsha, L. S. Cooper, E. Obarzanek, P. J. Elmer, V. J. Stevens, W. M. Vollmer, P. H. Lin, L. P. Svetkey, et al. 2003. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *Jama* 289:2083–93. doi:10.1001/jama.289.16.2083.

Appel, L. J., T. J. Moore, E. Obarzanek, W. M. Vollmer, L. P. Svetkey, F. M. Sacks, G. A. Bray, T. M. Vogt, J. A. Cutler, M. M. Windhauser, et al. 1997. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *The New England Journal of Medicine* 336:1117–24. doi:10.1056/NEJM199704173361601.

Ard, J. D., B. Gower, G. Hunter, C. S. Ritchie, D. L. Roth, A. Goss, B. C. Wingo, E. V. Bodner, C. J. Brown, D. Bryan, et al. 2016. Effects of calorie restriction in obese older adults: The CROSSROADS randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 73:73–80.

Azadbakht, L., P. Mirmiran, A. Esmailzadeh, T. Azizi, and F. Azizi. 2005. Beneficial effects of a Dietary Approaches to Stop Hypertension eating plan on features of the metabolic syndrome. *Diabetes Care* 28:2823–31.

Blumenthal, J. A., M. A. Babyak, A. Hinderliter, L. L. Watkins, L. Craighead, P. H. Lin, C. Caccia, J. Johnson, R. Waugh, and A. Sherwood. 2010. Effects of the DASH diet alone and in combination with exercise and weight loss on blood pressure and cardiovascular biomarkers in men and women with high blood pressure: the ENCORE study. *Archives of Internal Medicine* 170:126–35.

Brehm, B. J., B. L. Lattin, S. S. Summer, J. A. Boback, G. M. Gilchrist, R. J. Jandacek, and D. A. D'Alessio. 2009. One-year comparison of a high-monounsaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care* 32:215–20.

Brinkworth, G. D., M. Noakes, J. B. Keogh, N. D. Luscombe, G. A. Wittert, and P. M. Clifton. 2004a. Long-term effects of a high-protein, low-carbohydrate diet on weight control and cardiovascular risk markers in obese hyperinsulinemic subjects. *International Journal of Obesity and Related Metabolic Disorders* 28:661–70. doi:10.1038/sj.ijo.0802617.

Brinkworth, G. D., M. Noakes, B. Parker, P. Foster, and P. M. Clifton. 2004b. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia* 47:1677–86. doi:10.1007/s00125-004-1511-7.

Bucher, H. C., G. H. Guyatt, L. E. Griffith, and S. D. Walter. 1997. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *Journal of Clinical Epidemiology* 50:683–91.

Bueno, N. B., I. S. de Melo, S. L. de Oliveira, and T. da Rocha Ataíde. 2013. Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: a meta-analysis of randomised controlled trials. *British Journal of Nutrition* 110:1178–87.

Campbell, D. D., and K. A. Meckling. 2012. Effect of the protein:carbohydrate ratio in hypoenergetic diets on metabolic syndrome risk factors in exercising overweight and obese women. *British Journal of Nutrition* 108:1658–71.

Chaimani, A., D. M. Caldwell, T. Li, J. P. Higgins, and G. Salanti. 2017. Additional considerations are required when preparing a protocol for a systematic review with multiple interventions. *Journal of Clinical Epidemiology* 83:65–74.

Chaimani, A., J. P. Higgins, D. Mavridis, P. Spyridonos, and G. Salanti. 2013. Graphical tools for network meta-analysis in STATA. *Plos One* 8:e76654.

- Chaimani, A., and G. Salanti. 2015. Visualizing assumptions and results in network meta-analysis: The network graphs package. *Stata Journal* 15:905–50.
- Chalmers, J., T. Morgan, A. Doyle, B. Dickson, J. Hopper, J. Mathews, G. Matthews, R. Moulds, J. Myers, C. Nowson, et al. 1986. Australian National Health and Medical Research Council dietary salt study in mild hypertension. *Journal of Hypertension. Supplement* 4:S629–637.
- Clifton, P. M., M. Noakes, and J. B. Keogh. 2004. Very low-fat (12%) and high monounsaturated fat (35%) diets do not differentially affect abdominal fat loss in overweight, nondiabetic women. *The Journal of Nutrition* 134:1741–45.
- Coppell, K. J., M. Kataoka, S. M. Williams, A. W. Chisholm, S. M. Vorgers, and J. I. Mann. 2010. Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment—Lifestyle Over and Above Drugs in Diabetes (LOADD) study: randomised controlled trial. *Bmj (Clinical Research Ed.)* 341:c3337.
- Cornelissen, V. A., and N. A. Smart. 2013. Exercise training for blood pressure: a systematic review and meta-analysis. *Journal of the American Heart Association* 2:e004473.
- Costa, F. V., E. Ambrosioni, L. Montebugnoli, L. Paccaloni, L. Vasconi, and B. Magnani. 1981. Effects of a low-salt diet and of acute salt loading on blood pressure and intralymphocytic sodium concentration in young subjects with borderline hypertension. *Clinical Science (Lond)* 61 (Suppl 7):21s–3s.
- Dalle Grave, R., S. Calugi, I. Gavasso, M. El Ghoch, and G. Marchesini. 2013. A randomized trial of energy-restricted high-protein versus high-carbohydrate, low-fat diet in morbid obesity. *Obesity (Silver Spring)* 21:1774–81.
- Daly, M. E., R. Paisey, R. Paisey, B. A. Millward, C. Eccles, K. Williams, S. Hammersley, K. M. MacLeod, and T. J. Gale. 2006. Short-term effects of severe dietary carbohydrate-restriction advice in Type 2 diabetes—a randomized controlled trial. *Diabetic Medicine* 23:15–20.
- Dansinger, M. L., J. A. Gleason, J. L. Griffith, H. P. Selker, and E. J. Schaefer. 2005. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *Jama* 293:43–53.
- Delbridge, E. A., L. A. Prendergast, J. E. Pritchard, and J. Proietto. 2009. One-year weight maintenance after significant weight loss in healthy overweight and obese subjects: does diet composition matter? *The American Journal of Clinical Nutrition* 90:1203–14.
- Deluis, D. A., M. G. Sagrado, R. Aller, O. Izaola, and R. Conde. 2010. Effects of C358A missense polymorphism of the degrading enzyme fatty acid amide hydrolase on weight loss, adipocytokines, and insulin resistance after 2 hypocaloric diets. *Metabolism* 59:1387–92.
- Dias, S., N. J. Welton, D. M. Caldwell, and A. E. Ades. 2010. Checking consistency in mixed treatment comparison meta-analysis. *Stat Med* 29:932–44.
- Dodson, P. M., M. Beevers, R. Hallworth, M. J. Webberley, R. F. Fletcher, and K. G. Taylor. 1989. Sodium restriction and blood pressure in hypertensive type II diabetics: randomised blind controlled and cross-over studies of moderate sodium restriction and sodium supplementation. *Bmj (Clinical Research Ed.)* 298:227–30.
- Eckel, R. H., J. M. Jakicic, J. D. Ard, J. M. de Jesus, N. Houston Miller, V. S. Hubbard, I. M. Lee, A. H. Lichtenstein, C. M. Loria, B. E. Millen, et al. 2014. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 129:S76–99.
- Edwards, K. M., K. L. Wilson, J. Sadjia, M. G. Ziegler, and P. J. Mills. 2011. Effects on blood pressure and autonomic nervous system function of a 12-week exercise or exercise plus DASH-diet intervention in individuals with elevated blood pressure. *Acta Physiologica (Oxf)* 203:343–50.
- Elmer, P. J., E. Obarzanek, W. M. Vollmer, D. Simons-Morton, V. J. Stevens, D. R. Young, P. H. Lin, C. Champagne, D. W. Harsha, L. P. Svetkey, et al. 2006. Effects of comprehensive lifestyle modification on diet, weight, physical fitness, and blood pressure control: 18-month results of a randomized trial. *Annals of Internal Medicine* 144:485–95.
- Erwtaman, T. M., N. Nagelkerke, J. Lubsen, M. Koster, and A. J. Dunning. 1984. Beta blockade, diuretics, and salt restriction for the management of mild hypertension: a randomised double blind trial. *British Medical Journal (Clin Res Ed)* 289:406–409.
- Esposito, K., M. I. Maiorino, M. Ciotola, C. Di Palo, P. Scognamiglio, M. Gicchino, M. Petrizzo, F. Saccomanno, F. Beneduce, A. Ceriello, et al. 2009. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Annals of Internal Medicine* 151:306–14.
- Esposito, K., R. Marfella, M. Ciotola, C. Di Palo, F. Giugliano, G. Giugliano, M. D'Armiento, F. D'Andrea, and D. Giugliano. 2004. Effect of a mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *Jama* 292:1440–46.
- Ettehad, D., C. A. Emdin, A. Kiran, S. G. Anderson, T. Callender, J. Emberson, J. Chalmers, A. Rodgers, and K. Rahimi. 2016. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 387:957–67.
- Evans, C. E., D. C. Greenwood, D. E. Threapleton, C. P. Gale, C. L. Cleghorn, and V. J. Burley. 2017. Glycemic index, glycemic load, and blood pressure: a systematic review and meta-analysis of randomized controlled trials. *The American Journal of Clinical Nutrition* 105:1176–90.
- Fekete, A. A., D. I. Givens, and J. A. Lovegrove. 2015. Casein-derived lactotripeptides reduce systolic and diastolic blood pressure in a meta-analysis of randomised clinical trials. *Nutrients* 7:659–81.
- Frisch, S., A. Zittermann, H. K. Berthold, C. Gotting, J. Kuhn, K. Kleesiek, P. Stehle, and H. Kortke. 2009. A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *Cardiovasc Diabetol* 8:36.
- Gay, H. C., S. G. Rao, V. Vaccarino, and M. K. Ali. 2016. Effects of different dietary interventions on blood pressure: Systematic review and meta-analysis of randomized controlled trials. *Hypertension* 67:733–39.
- Gordon, N. F., C. B. Scott, and B. D. Levine. 1997. Comparison of single versus multiple lifestyle interventions: are the antihypertensive effects of exercise training and diet-induced weight loss additive? *American Journal of Cardiology* 79:763–67.
- Guldbrand, H., B. Dizdar, B. Bunjaku, T. Lindstrom, M. Bachrach-Lindstrom, M. Fredrikson, C. J. Ostgren, and F. H. Nystrom. 2012. In type 2 diabetes, randomisation to advice to follow a low-carbohydrate diet transiently improves glycaemic control compared with advice to follow a low-fat diet producing a similar weight loss. *Diabetologia* 55:2118–27.
- Haider, L. M., L. Schwingshackl, G. Hoffmann, and C. Ekmekcioglu. 2017. The effect of vegetarian diets on iron status in adults: A systematic review and meta-analysis. *Critical Reviews in Food Science and Nutrition* 0:1–16.
- Han, H., X. Fang, X. Wei, Y. Liu, Z. Jin, Q. Chen, Z. Fan, J. Aaseth, A. Hiyoshi, J. He, et al. 2017. Dose-response relationship between dietary magnesium intake, serum magnesium concentration and risk of hypertension: a systematic review and meta-analysis of prospective cohort studies. *Nutrition Journal* 16:26.
- Heilbronn, L. K., M. Noakes, and P. M. Clifton. 1999. Effect of energy restriction, weight loss, and diet composition on plasma lipids and glucose in patients with type 2 diabetes. *Diabetes Care* 22:889–95.
- Higgins, J. P., D. G. Altman, P. C. Gotzsche, P. Juni, D. Moher, A. D. Oxman, J. Savovic, K. F. Schulz, L. Weeks, and J. A. Sterne. 2011. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *Bmj (Clinical Research Ed.)* 343:d5928.
- Higgins, J. P., D. Jackson, J. K. Barrett, G. Lu, A. E. Ades, and I. R. White. 2012. Consistency and inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Research Synthesis Methods* 3:98–110.
- Higgins, J. P. T., and S. Green. 2011. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.
- Houston, M. C. 2011. The importance of potassium in managing hypertension. *Current Hypertension Reports* 13:309–17.
- Hutton, B., G. Salanti, D. M. Caldwell, A. Chaimani, C. H. Schmid, C. Cameron, J. P. Ioannidis, S. Straus, K. Thorlund, J. P. Jansen, et al. 2015. The PRISMA extension statement for reporting of systematic reviews incorporating

- network meta-analyses of health care interventions: checklist and explanations. *Annals of Internal Medicine* 162:777–84.
- Iqbal, N., M. L. Vetter, R. H. Moore, J. L. Chittams, C. V. Dalton-Bakes, M. Dowd, C. Williams-Smith, S. Cardillo, and T. A. Wadden. 2010. Effects of a low-intensity intervention that prescribed a low-carbohydrate vs. a low-fat diet in obese, diabetic participants. *Obesity (Silver Spring)* 18:1733–38.
- Itsiopoulos, C., L. Brazionis, M. Kaimakamis, M. Cameron, J. D. Best, K. O'Dea, and K. Rowley. 2011. Can the Mediterranean diet lower HbA1c in type 2 diabetes? Results from a randomized cross-over study. *Nutrition, Metabolism & Cardiovascular Diseases* 21:740–47.
- Jackson, D., J. K. Barrett, S. Rice, I. R. White, and J. P. Higgins. 2014. A design-by-treatment interaction model for network meta-analysis with random inconsistency effects. *Stat Med* 33:3639–54.
- Jonsson, T., Y. Granfeldt, B. Ahren, U. C. Branell, G. Palsson, A. Hansson, M. Soderstrom, and S. Lindeberg. 2009. Beneficial effects of a Paleolithic diet on cardiovascular risk factors in type 2 diabetes: a randomized cross-over pilot study. *Cardiovascular Diabetology* 8:35.
- Jula, A. M., and H. M. Karanko. 1994. Effects on left ventricular hypertrophy of long-term nonpharmacological treatment with sodium restriction in mild-to-moderate essential hypertension. *Circulation* 89:1023–31.
- Kim, H. H., Y. J. Kim, S. Y. Lee, D. W. Jeong, J. G. Lee, Y. H. Yi, Y. H. Cho, E. J. Choi, and H. J. Kim. 2014. Interactive effects of an isocaloric high-protein diet and resistance exercise on body composition, ghrelin, and metabolic and hormonal parameters in untrained young men: A randomized clinical trial. *Journal of Diabetes Investigation* 5:242–47.
- Kirpizidis, H., A. Stavratsi, and P. Geleris. 2005. Assessment of quality of life in a randomized clinical trial of candesartan only or in combination with DASH diet for hypertensive patients. *Journal of Cardiology* 46:177–82.
- Klemsdal, T. O., I. Holme, H. Nerland, T. R. Pedersen, and S. Tonstad. 2010. Effects of a low glycemic load diet versus a low-fat diet in subjects with and without the metabolic syndrome. *Nutrition, Metabolism & Cardiovascular Diseases* 20:195–201.
- Krahn, U., H. Binder, and J. Konig. 2013. A graphical tool for locating inconsistency in network meta-analyses. *BMC Medical Research Methodology* 13:35.
- Krebs, J. D., C. R. Elley, A. Parry-Strong, H. Lunt, P. L. Drury, D. A. Bell, E. Robinson, S. A. Moyes, and J. I. Mann. 2012. The Diabetes Excess Weight Loss (DEWL) Trial: a randomised controlled trial of high-protein versus high-carbohydrate diets over 2 years in type 2 diabetes. *Diabetologia* 55:905–14.
- Law, M., N. Wald, and J. Morris. 2003. Lowering blood pressure to prevent myocardial infarction and stroke: a new preventive strategy. *Health Technology Assessment* 7:1–94.
- Lawes, C. M., S. Vander Hoorn, and A. Rodgers. 2008. Global burden of blood-pressure-related disease, 2001. *Lancet* 371:1513–18.
- Lean, M. E., T. S. Han, T. Prvan, P. R. Richmond, and A. Avenell. 1997. Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *European Journal of Clinical Nutrition* 51:243–48.
- Leucht, S., A. Chaimani, A. S. Cipriani, J. M. Davis, T. A. Furukawa, and G. Salanti. 2016. Network meta-analyses should be the highest level of evidence in treatment guidelines. *European Archives of Psychiatry and Clinical Neuroscience* 266:477–80.
- Lewington, S., R. Clarke, N. Qizilbash, R. Peto, and R. Collins. 2002. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 360:1903–13.
- Lima, S. T., B. da Silva Nalin de Souza, A. K. Franca, N. Salgado Filho, and R. Sichieri. 2013. Dietary approach to hypertension based on low glycaemic index and principles of DASH (Dietary Approaches to Stop Hypertension): a randomised trial in a primary care service. *British Journal of Nutrition* 110:1472–79.
- Liu, X., G. Zhang, X. Ye, H. Li, X. Chen, L. Tang, Y. Feng, I. Shai, M. J. Stampfer, F. B. Hu, and X. Lin. 2013. Effects of a low-carbohydrate diet on weight loss and cardiometabolic profile in Chinese women: a randomised controlled feeding trial. *British Journal of Nutrition* 110:1444–53.
- Luger, M., B. Holstein, K. Schindler, R. Kruschitz, and B. Ludvik. 2013. Feasibility and efficacy of an isocaloric high-protein vs. standard diet on insulin requirement, body weight and metabolic parameters in patients with type 2 diabetes on insulin therapy. *Experimental and Clinical Endocrinology & Diabetes* 121:286–94.
- Ma, Y., B. C. Olendzki, P. A. Merriam, D. E. Chiriboga, A. L. Culver, W. Li, J. R. Hebert, I. S. Ockene, J. A. Griffith, and S. L. Pagoto. 2008. A randomized clinical trial comparing low-glycemic index versus ADA dietary education among individuals with type 2 diabetes. *Nutrition (Burbank, Los Angeles County, Calif.)* 24:45–56.
- Makela, P., T. Vahlberg, I. Kantola, R. Vesalainen, and A. Jula. 2008. The effects of a 6-month sodium restriction on cardiac autonomic function in patients with mild to moderate essential hypertension. *American Journal of Hypertension* 21:1183–87.
- Mancia, G., R. Fagard, K. Narkiewicz, J. Redon, A. Zanchetti, M. Bohm, T. Christiaens, R. Cifkova, G. De Backer, A. Dominiczak, et al. 2013a. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Journal of Hypertension* 31:1281–357.
- Mancia, G., R. Fagard, K. Narkiewicz, J. Redon, A. Zanchetti, M. Bohm, T. Christiaens, R. Cifkova, G. De Backer, A. Dominiczak, et al. 2013b. 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal* 34:2159–219.
- Manheimer, E. W., E. J. van Zuuren, Z. Fedorowicz, and H. Pijl. 2015. Paleolithic nutrition for metabolic syndrome: systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 102:922–32.
- Mansoor, N., K. J. Vinknes, M. B. Veierod, and K. Retterstol. 2016. Effects of low-carbohydrate diets v. low-fat diets on body weight and cardiovascular risk factors: a meta-analysis of randomised controlled trials. *British Journal of Nutrition* 115:466–79.
- Mavridis, D., M. Giannatsi, A. Cipriani, and G. Salanti. 2015. A primer on network meta-analysis with emphasis on mental health. *Evidence-Based Mental Health* 18:40–46.
- Mellberg, C., S. Sandberg, M. Ryberg, M. Eriksson, S. Brage, C. Larsson, T. Olsson, and B. Lindahl. 2014. Long-term effects of a Palaeolithic-type diet in obese postmenopausal women: a 2-year randomized trial. *European Journal of Clinical Nutrition* 68:350–57.
- NCD-Risk-Factor-Collaboration. 2017. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet* 389:37–55.
- Ndanuko, R. N., L. C. Tapsell, K. E. Charlton, E. P. Neale, and M. J. Batterham. 2016. Dietary patterns and blood pressure in adults: A systematic review and meta-analysis of randomized controlled trials. *Advances in Nutrition* 7:76–89.
- Neter, J. E., B. E. Stam, F. J. Kok, D. E. Grobbee, and J. M. Geleijnse. 2003. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 42:878–84.
- Nicholson, A. S., M. Sklar, N. D. Barnard, S. Gore, R. Sullivan, and S. Browning. 1999. Toward improved management of NIDDM: A randomized, controlled, pilot intervention using a lowfat, vegetarian diet. *Preventive Medicine* 29:87–91.
- Nissensohn, M., B. Roman-Vinas, A. Sanchez-Villegas, S. Piscopo, and L. Serra-Majem. 2016. The effect of the mediterranean diet on hypertension: A systematic review and meta-analysis. *Journal of Nutrition Education and Behavior* 48:42–53.e41.
- Nowson, C. A., A. Worsley, C. Margerison, M. K. Jorna, S. J. Godfrey, and A. Booth. 2005. Blood pressure change with weight loss is affected by diet type in men. *The American Journal of Clinical Nutrition* 81:983–89.
- Philippou, E., C. Bovill-Taylor, C. Rajkumar, M. L. Vampa, E. Ntatsaki, A. E. Brynes, M. Hickson, and G. S. Frost. 2009. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 58:1703–1708.
- Poulsen, S. K., C. Crone, A. Astrup, and T. M. Larsen. 2015. Long-term adherence to the New Nordic Diet and the effects on body weight, anthropometry and blood pressure: a 12-month follow-up study. *European Journal of Nutrition* 54:67–76.

- Riley, R. D., J. P. Higgins, and J. J. Deeks. 2011. Interpretation of random effects meta-analyses. *Bmj (Clinical Research Ed.)* 342:d549.
- Rock, C. L., S. W. Flatt, B. Pakiz, K. S. Taylor, A. F. Leone, K. Brelje, D. D. Heath, E. L. Quintana, and N. E. Sherwood. 2014. Weight loss, glycaemic control, and cardiovascular disease risk factors in response to differential diet composition in a weight loss program in type 2 diabetes: a randomized controlled trial. *Diabetes Care* 37:1573–80.
- Sacks, F. M., L. P. Svetkey, W. M. Vollmer, L. J. Appel, G. A. Bray, D. Harsha, E. Obarzanek, P. R. Conlin, E. R. Miller, 3rd, D. G. Simons-Morton, et al. 2001. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *The New England Journal of Medicine* 344:3–10.
- Salanti, G. 2012. Indirect and mixed-treatment comparison, network, or multiple-treatments meta-analysis: many names, many benefits, many concerns for the next generation evidence synthesis tool. *Research Synthesis Methods* 3:80–97.
- Salanti, G., A. E. Ades, and J. P. Ioannidis. 2011. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of Clinical Epidemiology* 64:163–71.
- Salanti, G., C. Del Giovane, A. Chaimani, D. M. Caldwell, and J. P. Higgins. 2014. Evaluating the quality of evidence from a network meta-analysis. *Plos One* 9:e99682.
- Saneei, P., A. Salehi-Abargouei, A. Esmailzadeh, and L. Azadbakht. 2014. Influence of Dietary Approaches to Stop Hypertension (DASH) diet on blood pressure: a systematic review and meta-analysis on randomized controlled trials. *Nutrition, Metabolism & Cardiovascular Diseases* 24:1253–61.
- Santesso, N., E. A. Akl, M. Bianchi, A. Mente, R. Heels-Ansdell, D., and H. J. Schunemann. 2012. Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *European Journal of Clinical Nutrition* 66:780–88.
- Schwingshackl, L., A. Chaimani, G. Hoffmann, C. Schwedhelm, and H. Boeing. 2017a. Impact of different dietary approaches on blood pressure in hypertensive and prehypertensive patients: protocol for a systematic review and network meta-analysis. *BMJ Open* 7:e014736.
- Schwingshackl, L., A. Chaimani, G. Hoffmann, C. Schwedhelm, and H. Boeing. 2018. A network meta-analysis on the comparative efficacy of different dietary approaches on glycaemic control in patients with type 2 diabetes mellitus. *European Journal of Epidemiology* 33:157–70.
- Schwingshackl, L., S. Dias, B. Strasser, and G. Hoffmann. 2013. Impact of different training modalities on anthropometric and metabolic characteristics in overweight/obese subjects: a systematic review and network meta-analysis. *Plos One* 8:e82853.
- Schwingshackl, L., L. P. Hobl, and G. Hoffmann. 2015a. Effects of low glycaemic index/low glycaemic load vs. high glycaemic index/ high glycaemic load diets on overweight/obesity and associated risk factors in children and adolescents: a systematic review and meta-analysis. *Nutrition Journal* 14:87.
- Schwingshackl, L., and G. Hoffmann. 2013a. Comparison of effects of long-term low-fat vs high-fat diets on blood lipid levels in overweight or obese patients: a systematic review and meta-analysis. *Journal of the Academy of Nutrition and Dietetics* 113:1640–61.
- Schwingshackl, L., and G. Hoffmann. 2013b. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutrition Journal* 12:48.
- Schwingshackl, L., and G. Hoffmann. 2013c. Long-term effects of low glycaemic index/load vs. high glycaemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. *Nutr Metab Cardiovasc Dis* 23:699–706.
- Schwingshackl, L., and G. Hoffmann. 2014a. Adherence to Mediterranean diet and risk of cancer: a systematic review and meta-analysis of observational studies. *Int J Cancer* 135:1884–97.
- Schwingshackl, L., and G. Hoffmann. 2014b. Comparison of the long-term effects of high-fat v. low-fat diet consumption on cardiometabolic risk factors in subjects with abnormal glucose metabolism: a systematic review and meta-analysis. *British Journal of Nutrition* 111:2047–58.
- Schwingshackl, L., and G. Hoffmann. 2014c. Mediterranean dietary pattern, inflammation and endothelial function: a systematic review and meta-analysis of intervention trials. *Nutrition, Metabolism & Cardiovascular Diseases* 24:929–39.
- Schwingshackl, L., and G. Hoffmann. 2015a. Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis of observational studies. *Cancer Medicine* 4:1933–47.
- Schwingshackl, L., and G. Hoffmann. 2015b. Diet quality as assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension score, and health outcomes: a systematic review and meta-analysis of cohort studies. *Journal of the Academy of Nutrition and Dietetics* 115:780–800.e785.
- Schwingshackl, L., and G. Hoffmann. 2016. Does a Mediterranean-type diet reduce cancer risk? *Current Nutrition Reports* 5:9–17.
- Schwingshackl, L., B. Missbach, S. Dias, J. Konig, and G. Hoffmann. 2014. Impact of different training modalities on glycaemic control and blood lipids in patients with type 2 diabetes: a systematic review and network meta-analysis. *Diabetologia* 57:1789–97.
- Schwingshackl, L., B. Missbach, J. Konig, and G. Hoffmann. 2015b. Adherence to a Mediterranean diet and risk of diabetes: a systematic review and meta-analysis. *Public Health Nutr.* 18:1292–99.
- Schwingshackl, L., C. Schwedhelm, G. Hoffmann, S. Knuppel, K. Iqbal, V. Andriolo, A. Bechthold, S. Schlesinger, and H. Boeing. 2017b. Food groups and risk of hypertension: A systematic review and dose-response meta-analysis of prospective studies. *Advances in Nutrition* 8:793–803.
- Schwingshackl, L., B. Strasser, and G. Hoffmann. 2011. Effects of monounsaturated fatty acids on cardiovascular risk factors: a systematic review and meta-analysis. *Annals of Nutrition and Metabolism* 59:176–86.
- Shai, I., D. Schwarzfuchs, Y. Henkin, D. R. Shahar, S. Witkow, I. Greenberg, R. Golan, D. Fraser, A. Bolotin, H. Vardi, et al. 2008. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *The New England Journal of Medicine* 359:229–41.
- Siervo, M., J. Lara, S. Chowdhury, A. Ashor, C. Oggioni, and J. C. Mathers. 2015. Effects of the Dietary Approach to Stop Hypertension (DASH) diet on cardiovascular risk factors: a systematic review and meta-analysis. *British Journal of Nutrition* 113:1–15.
- Silman, A. J., C. Locke, P. Mitchell, and P. Humpherson. 1983. Evaluation of the effectiveness of a low sodium diet in the treatment of mild to moderate hypertension. *Lancet*. 1:1179–82.
- Smith, S. C. Jr., E. J. Benjamin, R. O. Bonow, L. T. Braun, M. A. Creager, B. A. Franklin, R. J. Gibbons, S. M. Grundy, L. F. Hiratzka, D. W. Jones, et al. 2011. AHA/ACC secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease: 2011 update: a guideline from the American heart association and American college of cardiology foundation. *Circulation* 124:2458–73.
- StataCorp. 2015. *Stata statistical software: Release 14*. College Station, TX: StataCorp LP.
- Stern, L., N. Iqbal, P. Seshadri, K. L. Chicano, D. A. Daily, J. McGrory, M. Williams, E. J. Gracely, and F. F. Samaha. 2004. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Annals of Internal Medicine* 140:778–85.
- Tang, M., C. L. Armstrong, H. J. Leidy, and W. W. Campbell. 2013. Normal vs. high-protein weight loss diets in men: effects on body composition and indices of metabolic syndrome. *Obesity (Silver Spring)*. 21:E204–210.
- Tay, J., G. D. Brinkworth, M. Noakes, J. Keogh, and P. M. Clifton. 2008. Metabolic effects of weight loss on a very-low-carbohydrate diet compared with an isocaloric high-carbohydrate diet in abdominally obese subjects. *Journal of the American College of Cardiology* 51:59–67.
- Tay, J., N. D. Luscombe-Marsh, C. H. Thompson, M. Noakes, J. D. Buckley, G. A. Wittert, W. S. Jr Yancy, and G. D. Brinkworth. 2015. Comparison of low- and high-carbohydrate diets for type 2 diabetes management: a randomized trial. *The American Journal of Clinical Nutrition* 102:780–90.
- Toledo, E., F. B. Hu, R. Estruch, and e. a. 2013. Effect of the Mediterranean diet on blood pressure in the PREDIMED trial: results from a randomized controlled trial. *BMC Medicine* 11:207.
- Trichopoulou, A., T. Costacou, C. Bamia, and D. Trichopoulos. 2003. Adherence to a Mediterranean diet and survival in a Greek population. *The New England Journal of Medicine* 348:2599–608.

- Uusitupa, M., K. Hermansen, M. J. Savolainen, U. Schwab, M. Kolehmainen, L. Brader, L. S. Mortensen, L. Cloetens, A. Johansson-Persson, G. Onning, et al. 2013. Effects of an isocaloric healthy Nordic diet on insulin sensitivity, lipid profile and inflammation markers in metabolic syndrome – a randomized study (SYSDIET). *Journal of Internal Medicine* 274:52–66.
- Uusitupa, M., J. Laitinen, O. Siitonen, E. Vanninen, and K. Pyorala. 1993. The maintenance of improved metabolic control after intensified diet therapy in recent type 2 diabetes. *Diabetes Research and Clinical Practice* 19:227–38.
- Veum, V. L., J. Laupsa-Borge, O. Eng, E. Rostrup, T. H. Larsen, J. E. Nordrehaug, O. K. Nygard, J. V. Sagen, O. A. Gudbrandsen, S. N. Dankel, et al. 2017. Visceral adiposity and metabolic syndrome after very high-fat and low-fat isocaloric diets: a randomized controlled trial. *The American Journal of Clinical Nutrition* 105:85–99.
- von Haehling, S., K. Stellos, N. Qusar, M. Gawaz, and B. Bigalke. 2013. Weight reduction in patients with coronary artery disease: comparison of Traditional Tibetan Medicine and Western diet. *International Journal of Cardiology* 168:1509–15.
- Walker, K. Z., K. O'Dea, G. C. Nicholson, and J. G. Muir. 1995. Dietary composition, body weight, and NIDDM. Comparison of high-fiber, high-carbohydrate, and modified-fat diets. *Diabetes Care* 18:401–403.
- Watkins, L. L., A. Sherwood, M. Feinglos, A. Hinderliter, M. Babyak, E. Gullette, R. Waugh, and J. A. Blumenthal. 2003. Effects of exercise and weight loss on cardiac risk factors associated with syndrome X. *Archives of Internal Medicine* 163:1889–95.
- Watson, N., K. Dyer, J. Buckley, G. Brinkworth, A. Coates, G. Parfitt, P. Howe, M. Noakes, and K. Murphy. 2016. Effects of low-fat diets differing in protein and carbohydrate content on cardiometabolic risk factors during weight loss and weight maintenance in obese adults with Type 2 Diabetes. *Nutrients* 8(5):289.
- Westman, E. C., W. S. Jr. Yancy, J. C. Mavropoulos, M. Marquart, and J. R. McDuffie. 2008. The effect of a low-carbohydrate, ketogenic diet versus a low-glycemic index diet on glycemic control in type 2 diabetes mellitus. *Nutr Metab (Lond)* 5:36.
- Whelton, P., e. a. 1997. Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Archives of Internal Medicine* 157:657–67.
- Whelton, P. K., L. J. Appel, M. A. Espeland, W. B. Applegate, W. H. Jr. Ettinger, J. B. Kostis, S. Kumanyika, C. R. Lacy, K. C. Johnson, S. Folmar, et al. 1998. Sodium reduction and weight loss in the treatment of hypertension in older persons: a randomized controlled trial of non-pharmacologic interventions in the elderly (TONE). TONE Collaborative Research Group. *Jama*. 279:839–46.
- Whelton, S. P., A. D. Hyre, B. Pedersen, Y. Yi, P. K. Whelton, and J. He. 2005. Effect of dietary fiber intake on blood pressure: a meta-analysis of randomized, controlled clinical trials. *Journal of Hypertension* 23:475–81.
- White, I. R. 2015. Network meta-analysis. *Stata Journal* 15:951–85.
- World-Health-Organization. 2012. Raised blood pressure: Situation and trends. *Global Health Observatory* http://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence/en/. (Accessed 04.02.2016).
- Wycherley, T. P., L. J. Moran, P. M. Clifton, M. Noakes, and G. D. Brinkworth. 2012. Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *The American Journal of Clinical Nutrition* 96:1281–98.
- Wycherley, T. P., M. Noakes, P. M. Clifton, X. Cleanthous, J. B. Keogh, and G. D. Brinkworth. 2010. A high-protein diet with resistance exercise training improves weight loss and body composition in overweight and obese patients with type 2 diabetes. *Diabetes Care* 33:969–76.
- Yokoyama, Y., K. Nishimura, N. D. Barnard, M. Takegami, M. Watanabe, A. Sekikawa, T. Okamura, and Y. Miyamoto. 2014. Vegetarian diets and blood pressure: a meta-analysis. *JAMA Internal Medicine* 174: 577–87.